

INVESTIGATING THE ASSOCIATION BETWEEN CYBERCHONDRIA AND HEALTH ANXIETY

Sharvari S Gupte^{1*}

^{1*}Assistant Professor, Department of Psychology, KET's V. G. Vaze College of Arts, Science and Commerce (Autonomous), Mulund. (+91 9321382331, sharvarigupte261200@gmail.com)

*Corresponding Author

*Email: sharvarigupte261200@gmail.com

Abstract: This research explores a growing health concern of the digital era—cyberchondria—and its link to anxiety among young adults aged 18-25 in Mumbai, India. Utilizing a mixed-method approach, the study includes a quantitative survey of 72 participants (N=72) aged 18-24, alongside qualitative interviews with experienced physicians. The quantitative analysis, conducted using Pearson's Product-Moment Correlation, revealed a strong positive correlation between cyberchondria and anxiety ($r=0.76$, $p<0.001$). Qualitative insights underscored the challenges faced by doctors in managing patients who are digitally informed, stressing the need for enhanced digital literacy and improved communication strategies. This study addresses a crucial gap in the existing literature by focusing on the Indian context, offering practical recommendations such as the introduction of digital health literacy programs and evidence-based interventions to mitigate health-related anxieties. However, limitations include the use of self-reported data and a narrow demographic focus, suggesting a need for further research leveraging advanced technologies like AI to gain a deeper understanding of cyberchondria in the Indian population.

Keywords: Cyberchondria, Anxiety, Healthcare Professionals, Mental Health, Digital Age

Introduction

The digital era has dramatically expanded access to health information online, enabling individuals to take a proactive role in managing their healthcare. However, this increased access has also led to the rise of cyberchondria—excessive health anxiety triggered by online searches for medical information. Although extensive research on cyberchondria exists globally, there is a noticeable gap in studies focusing on its impact within the Indian population. This study aims to bridge this gap by examining cultural differences, technological literacy, and evolving healthcare landscapes specific to India. Understanding the perspectives of healthcare professionals, who have in-depth experience with cyberchondria, is essential for effectively managing patients affected by this condition.

Most existing research on cyberchondria is rooted in Western contexts, often overlooking the socio-cultural diversity unique to India. Current studies do not adequately capture how cultural nuances and varying levels of internet literacy influence the prevalence of cyberchondria in the Indian context. Furthermore, the role of healthcare providers in addressing this issue remains underexplored.

In today's digital age, the internet serves as a crucial repository of health information, empowering individuals to make informed decisions about their physical and mental well-being. However, the overwhelming availability of online health data has contributed to the phenomenon of cyberchondria, where excessive information consumption heightens health-related anxiety. This literature review examines key themes, conceptual frameworks, and empirical findings on cyberchondria, highlighting its prevalence, contributing factors, and its impact on mental health.

Literature Review

The term "cyberchondria" is a blend of "cyber" and "hypochondria," referring to a modern health concern driven by online searches for medical information. This concept was first introduced by Starcevic and Berle (2013) to describe a repetitive pattern of anxiety resulting from excessive online health-related searches, where perceived symptoms are linked to potential health problems. Their model highlights the compulsive nature of reassurance-seeking behaviors, which can amplify perceived health issues.

Cyberchondria prevalence rates differ across various demographic groups. McElroy and Shevlin (2019) found that younger individuals, females, and those with heightened health concerns are particularly susceptible to cyberchondria. However, the existing body of literature lacks a comprehensive understanding of how cultural differences and digital literacy influence the prevalence of cyberchondria. Key factors that contribute to the onset and persistence of this condition include information overload, confirmation bias, and availability heuristics, as noted by White and Horvitz (2009).

Norr and Albanese (2015) identified a bidirectional relationship between cyberchondria, anxiety, and stress, where each element reinforces the other. Concerns have been raised regarding the potential impact of cyberchondria on actual health behaviors. For instance, individuals may either avoid seeking medical advice due to fear of a serious diagnosis or engage in inappropriate self-treatment practices based on online information. Understanding effective strategies for managing cyberchondria is crucial to mitigate its negative effects.

Interventions focused on digital health literacy and promoting a balanced approach to online health information could be beneficial, as suggested by Muse et al. (2012). Cognitive Behavioral Therapy (CBT) has shown promise in addressing the cognitive distortions associated with cyberchondria (Fergus et al., 2018). This literature review explores the changing dynamics of cyberchondria, examining its prevalence, clinical mechanisms, and its effects on mental health.

While previous research has provided valuable insights, there remains a clear need for studies incorporating cultural diversity and a multidisciplinary approach, especially regarding the role of healthcare professionals in managing cyberchondria. Given the rapid growth in digital health, further research is essential to develop targeted interventions that foster a healthier relationship between individuals, online health resources, and mental well-being.

Research Methodology

This extensive study employs a mixed-methods approach to explore the complex relationship between cyberchondria and anxiety among the Indian population, while also gaining insights from healthcare professionals. The quantitative component involves the development and administration of a tailored questionnaire to assess the frequency, causes, and impact of cyberchondria on individuals' lives. Although standardized instruments like the Cyberchondria Severity Scale (CSS) informed the survey design, the actual scale was not directly administered. Demographic information (e.g., age, gender, education level) was gathered using a structured questionnaire.

The inclusion criteria for the survey include participants aged 18-24, with a basic understanding of English, access to the internet, and ownership of a mobile phone. Exclusion criteria consist of individuals with terminal illnesses, diagnosed mental health conditions, or those outside the specified age range. Convenience sampling was utilized to minimize random variation, and participants provided consent before completing the survey via Google Forms, an online platform. The study population comprises young adults aged 18-25 residing in Mumbai. Data analysis will involve inferential statistics, specifically Pearson's Product-Moment Correlation, to determine the association between cyberchondria and anxiety within the target demographic, thereby offering a quantitative perspective on health anxieties in the digital age in India.

The qualitative aspect of the study includes semi-structured interviews with healthcare professionals, particularly doctors, to gain in-depth insights into their perspectives on cyberchondria. The interviews aim to explore two key areas: the impact of cyberchondria on doctor-patient interactions and recommendations for effective healthcare interventions. Purposive sampling was employed to ensure a diverse range of medical specializations and experience levels, with inclusion criteria requiring doctors to have a minimum of seven years of experience in their respective fields. Qualitative data will be collected through interviews and analyzed thematically based on responses from healthcare professionals.

This study aims to provide a comprehensive understanding of the challenges cyberchondria poses within the Indian healthcare context, addressing existing research gaps and contributing to the development of future healthcare strategies in the digital era.

Results

The aim of this study was to explore the relationship between cyberchondria and anxiety, along with evaluation of healthcare professionals' perspective on the rising prevalence of cyberchondria in the Indian population. The population for this study was individuals belonging to the age group of 18-24 years residing in the city of Mumbai. A sample size (N) for the quantitative analysis was 72 respondents (N=72) consisting of 36 males and 36 females, and the survey was administered through google forms. For the qualitative analysis, 10 doctors from different specialisations were consulted. The hypothesis of the study was: There is a positive correlation between cyberchondria and anxiety levels in the Indian population, indicating that individuals who engage in excessive online health-related searches are more likely to experience higher levels of anxiety. Descriptive statistics was done to establish the correlation of the variables. Pearson Product Moment correlation was used to understand the relationship between the two variables.

Table 1.1 Descriptive statistics of Cyberchondria and Anxiety

	N	Mean	SD	Skewness	Kurtosis
Cyberchondria	72	16.287	5.574	0.041	-0.429
Anxiety	72	10.178	3.522	0.213	-0.391

Table 1.1 describes the variables of the study. The mean of the variable cyberchondria was 16.287 and the SD was 5.574. The mean of the variable anxiety was 10.178 and the SD was 3.522. The variables of Cyberchondria and anxiety were analysed for skewness and kurtosis and they were found to be in the acceptable range of skewness and kurtosis, as seen in the table 1.1.

Table 1.2 Pearson Product-Moment correlation between Cyberchondria and Anxiety

Pearson's Correlations

Variable		Column 2	Column 1
Cyberchondria	Pearson's r	—	
	p-value	—	
Anxiety	Pearson's r	0.755	—
	p-value	< .001	—

Table 1.2 presents the correlation findings among the variables: cyberchondria and anxiety. A strong and positive correlation of 0.76 was found between cyberchondria and anxiety levels in the sample of 72 respondents. The r value (0.76) is significant at $p < 0.01$ level. Hence, the results found are statistically significant and in line with the stated hypothesis.

Discussion

The term cyberchondria was popularised in the mid-1900s, by the British media by combining the words, cyber and hypochondria. Eventually cyberchondria became frequently used in academic literature and was used to describe individuals who excessively searched health related information, online (Loos, 2013). Cyberchondria as an illness is not mentioned in the DSM-V, but it is indirectly referenced as a diagnostic feature, describing Illness Anxiety Disorder as "excessively researching suspected illness on the internet."

Cyberchondria is not just related to searching for health information related to physical symptoms, but also related to the search of mental health information as well (Mathes et al., 2018). Researchers acknowledge that using the internet to look up health information does have advantages of informing individuals about the origin, prevention and management or the characteristics of certain health issues. However, when people feel worried or anxious about their health, they may turn to the internet to self-diagnose or to find assurance which might prove to be harmful as the resources on the internet might not always give them the correct diagnosis, or the internet has tendency to actually exaggerate the minimal symptoms that the individual might be experiencing.

According to McElroy & Shevlin (2014), cyberchondria consists of five dimensions, namely:

(1) Compulsion, describing how excessive searching for health information on the internet can hinder other activities; (2) Distress, reflecting the more subjective and deeper feelings associated with searching for health information on the internet; (3) Excessiveness, describing excessive and repeated searching for health information on the internet; (4) Seeking Finding Reassurance, increasing anxiety and the need to consult with professionals about information obtained from the internet; (5) Mistrust of Medical Professionals, the emergence of distrust towards professionals. As with every other area of human life, digital technology has transformed how humans, especially the young generation with their digital competence approaches healthcare. With an ocean of information available at their fingertips, youngsters tend to rely on the internet for any health-related information gathering, self-diagnosis and even seeking medical advice through online platforms. This has been observed to create a strain in the doctor-patient relationship as the patients visit the doctor with a preformed notion about their health issue and want the doctor to just confirm that and resist any deviant analysis.

It has been researched that people get anxious due to the fear of the unknown and uncertainty about an illness and heightens due to previous experiences faced while seeking help at the healthcare facilities and perceived lack of control. Shedding a light on their emotional needs is necessary for the provision of holistic care and minimising the adverse effects that anxiety can have on the treatment plans.

Anxiety and worry are two of the most prominent emotions experienced by patients across various healthcare contexts. It emanates from the apprehension surrounding the illness, fear of the unknown, and the potential implication of the diagnosis and the subsequent treatment plan. These anxieties are often amplified by factors such as perceived lack of control, cultural beliefs, and previous healthcare experiences. Healthcare professionals are rampantly discouraging and even warning the patients against using the internet to read about their symptoms or self-diagnose themselves. There is a need to inculcate the holistic approach in healthcare, and this will only be achieved by addressing the emotional needs of the patients and mitigating the adverse effects of anxiety on the patient's well-being

The themes which are identified through the narratives in doctor-patient interactions highlight the interplay between the advances in technology, emotional well-being, trust in the healthcare delivery system, and patient empowerment. As the healthcare environment in India continually changes, there is a need for all the stakeholders to pick up and adjust the dynamics and adapt more people-centred approaches. This includes approaches that place more emphasis on the attributes of communication, empathy, and collaboration, among others, and which ensure that health outcomes of the patient are maximal. There is a need for future research to build on these and develop more themes and strategies to improve and promote patient-doctor interactions and relationships in modern settings.

Conclusion

This research explores the complex relationship between cyberchondria and anxiety among young adults in Mumbai, India, aiming to illuminate how excessive online health information seeking influences mental well-being. By employing a mixed-method approach, the study offers a thorough understanding of both the prevalence of cyberchondria and healthcare professionals' perspectives on managing this emerging digital-age concern.

The quantitative analysis reveals a strong positive correlation between cyberchondria and anxiety, indicating that frequent online health searches are closely linked to heightened anxiety levels. On the qualitative side, insights from healthcare practitioners highlight the daily challenges of addressing concerns from digitally informed patients, emphasizing the urgent need for enhanced digital health literacy and improved communication strategies.

This research addresses a critical gap in existing literature by focusing on the distinct socio-cultural factors influencing cyberchondria in India, a region often overlooked in global studies. The findings suggest practical applications such as integrating digital health literacy into educational curricula and developing evidence-based strategies to manage health anxieties. These measures can contribute to fostering a more informed, resilient, and adaptable society.

However, the study acknowledges certain limitations, including its reliance on self-reported data and a specific demographic focus. It underscores the need for further research, particularly incorporating advanced technologies like Artificial Intelligence (AI), to deepen our understanding of cyberchondria in the evolving digital landscape of India.

In conclusion, this study provides valuable insights into the link between cyberchondria and anxiety in the Indian context, paving the way for more targeted and effective healthcare communication and intervention strategies. It highlights the importance of addressing digital health anxieties holistically, ensuring that both individuals and healthcare systems are better prepared to navigate the challenges of the digital era.

References

1. Starcevic, V., & Berle, D. (2013). Cyberchondria: towards a better understanding of excessive health-related Internet use. *Expert review of neurotherapeutics*, 13(2), 205–213. <https://doi.org/10.1586/ern.12.162>
2. McElroy, E., Kearney, M., Touhey, J., Evans, J., Cooke, Y., & Shevlin, M. (2019). The CSS-12: Development and Validation of a Short-Form Version of the Cyberchondria Severity Scale. *Cyberpsychology, behavior and social networking*, 22(5), 330–335. <https://doi.org/10.1089/cyber.2018.0624>
3. Ryen W. White and Eric Horvitz. 2009. Cyberchondria: Studies of the escalation of medical concerns in Web search. *ACM Trans. Inf. Syst.* 27, 4, Article 23 (November 2009), 37 pages. <https://doi.org/10.1145/1629096.1629101>
4. Norr, A. M., Albanese, B. J., Oglesby, M. E., Allan, N. P., & Schmidt, N. B. (2015). Anxiety sensitivity and intolerance of uncertainty as potential risk factors for cyberchondria. *Journal of Affective Disorders*, 174, 64-69. <https://doi.org/10.1016/j.jad.2014.11.023>.
5. Muse, K., McManus, F., Leung, C., Meghreblian, B., & Williams, J. M. G. (2012). Cyberchondriasis: fact or fiction? A preliminary examination of the relationship between health anxiety and searching for health information on the Internet. *Journal of anxiety disorders*, 26(1), 189-196. <https://doi.org/10.1016/j.janxdis.2011.11.005>.

6. Fergus, T. A., & Spada, M. M. (2018). Moving toward a metacognitive conceptualization of cyberchondria: Examining the contribution of metacognitive beliefs, beliefs about rituals, and stop signals. *Journal of anxiety disorders*, 60, 11-19. <https://doi.org/10.1016/j.janxdis.2018.09.003>.
7. McElroy, E., & Shevlin, M. (2014). The development and initial validation of the cyberchondria severity scale (CSS). *Journal of anxiety disorders*, 28(2), 259-265. <https://doi.org/10.1016/j.janxdis.2013.12.007>
8. Loos, A. (2013). Cyberchondria: Too Much Information for the Health Anxious Patient? *Journal of Consumer Health on the Internet*, 17(4), 439–445. <https://doi.org/10.1080/15398285.2013.833452>
9. Mathes, B.M., Norr, A.M., Allan, N.P., Albanese, B.J., Schmidt, N.B., 2018. Cyberchondria: overlap with health anxiety and unique relations with impairment, quality of life, and service utilization. *Psychiatry Res.* 261, 204–211.
10. <https://doi.org/10.1016/j.psychres.2018.01.002>