



FUNCTIONAL AND RADIOLOGICAL ANALYSIS OF MIDSHAFT CLAVICLE FRACTURE TREATED WITH TENS NAILING

Amit Joshi¹, Shrikrushna B.Shinde^{2*}, Dr Yash Haresh Ratanpal³, Dr. Anubhav Vichitra⁴, Vinayaka Reddy K C⁵, Dr Rizwan Yusuf Halai⁶

¹M. S Orthopaedics, Professor and Head, Department of orthopaedic, HBT Trauma Care Centre, Jogeshwari, ORCID iD: <https://orcid.org/0009-0003-7119-7656>, Email ID of author: Dr.amitjoshi85@gmail.com

^{2*}M. S Orthopaedics, Post MS Orthopaedic Resident Doctor, HBT Trauma care centre jogeshwari, Mumbai, Contact: +91-8390965313, E-mail: shindekrushna148@gmail.com, ORCID iD: <https://orcid.org/0009-0000-2478-1727>

³MS Ortho, Assistant Professor, Orthopaedics, 400101, Mumbai India, 0009-0002-8711-8193, Email ID yashratanpal89@gmail.com

⁴MS ORTHOPEDICS, Senior Resident, ORTHOPAEDICS, HBT TRAUMA CARE, Delhi University, 400037 MUMBAI INDIA, ORCID ID 0009-0007-9691-5123, Email ID ortho.anubhavvichitra.2019@gmail.com

⁵MS ORTHO, Senior Resident, ORTHOPAEDICS, HBT TRAUMA CARE, 400060, MUMBAI INDIA, ORCID ID 0009-0000-8915-437X, Email ID kevreddy15@gmail.com

⁶MS ORTHO, Senior Resident, ORTHOPAEDICS, HBT TRAUMA CARE, MUHS Nashik, 400037, MUMBAI INDIA, ORCID ID 0009-0008-1172-6700, Email ID rizwanhalai@yahoo.com

Received:- 19/07/2026

Revised:-21/08/2026

Accepted:-31/08/2026

Published:- 09/09/2026

Abstract

Introduction: Clavicle fracture is the most frequently and easily fractured bone due to fall on outstretched hand or direct impact. Mainly the midshaft clavicle fracture is 80% compare to medial and lateral end clavicle fracture. Amongst all fractures the incidence of clavicle fracture is 3-5%. Treatment consist of mainly two procedure open reduction with plate or minimally invasive close reduction with TENS nail.

Material & Methods: A prospective study was conducted from November 2022 to January 2025 with 30 patients whose plain radiographs shows midshaft clavicle fractures. This study was conducted at our orthopaedic institute which a tertiary care centre. Patients treated with TENS nailing and Functional outcomes were assessed using the Constant-Murley score, and radiological healing assessed with help of follow-up X-rays. Outcomes were analyzed at 4 weeks 8 weeks, 12 weeks, and 6 months post-operation. Implant removal done within 1 year and after 6 month of surgery

Results: Among 30 patients we achieved closed reduction of fracture in 25 patients with lesser blood loss while in remaining 5 patients required open reduction of fracture site using bone holding and artery forcep. In all patient clinical union occurs at 5 weeks on an average and radiologically union occurs on an average 10 weeks.

Conclusion: TENS nailing may offer a promising option for the treatment of clavicle shaft fractures as it is minimally invasive require less time for surgery (approx 30min) and patient can mobilize early with short span of hospital stay (1 day postoperatively). providing functional recovery with minimal complications and having superior cosmetic results.

Keywords: Midshaft Clavicle fracture, Titanium Elastic Nailing, Functional outcome, Radiological outcome, Prospective study, Fracture healing, Constant and Murley scoring

Introduction

Clavicle is the most frequently fracture bone either due to direct impact or fall on outstretched hand. among all fracture the incidence of clavicle fracture is 3-5%. various treatment modality available for clavicle fracture (1-8). there are mainly two operative approach which commonly in practice consist of open reduction using plate and close reduction using minimally invasive TENS nailing (1,5,8). although plating is standard technique but it has some disadvantages like non-union, large scar, infection and difficulty in application and removal. whereas, minimally invasive intramedullary TENS nailing have their own advantages and disadvantages. Titanium Elastic Nail System (TENS) because of less complication, rapid union rate, easy insertion and removal, small scar i.e. cosmetically superior (6,14)

Material & Methods

Study design : Prospective study.

Place of study: HBT Trauma care centre, Jogeshwari

Mumbai, Maharashtra a Tertiary care hospital

Duration of study: 25 Months

Sample size: 30

Inclusion Criteria:

- Adults aged >18 years of age
- OTA type A&B
- Displaced closed clavicle fracture or comminuted clavicle fractures.
- Midshaft clavicle fracture
- Bilateral clavicle fracture
- Clavicle fracture associated with scapula fracture

Exclusion Criteria:

- Open fractures i.e. compound fracture
- Pathological fractures
- Age <18 years
- Patients with comorbidities affecting fracture healing (e.g., diabetes, smoking).
- Medial and lateral end clavicle fracture

Surgical Technique

Operative procedure: After administration of General Anaesthesia, the patient will be placed in supine position with injured extremity prepared with 3 stage painting (savlon-spirit-betadine) and draped from the midline to the upper arm. Care will be taken to make sure that the sternoclavicular joint will be exposed well to facilitate easy entry point. Preoperatively, the shoulder region will be screened using image intensifier to confirm fracture site. A vertical skin incision will be made just lateral to the sternoclavicular joint. The subcutaneous fat will be incised. The pectoral fascia will be divided in line with the skin incision followed by careful elevation of the underlying musculature from the clavicle. The entry point will be then made using the awl and appropriate sized TENS nail will be inserted (The size of the nail will be measured using this formula = $0.4 \times \text{canal diameter in mm}$) on an average 2.5mm size used. Attempt will be made to close reduce the fracture. If the fracture could not be reduced by closed means, then a separate vertical incision will be used at the fracture site to achieve fracture reduction. Vertical incision will be used to open fracture site as it is parallel to Langer's lines and minimized the risk of damage to supraclavicular nerves to avoid dysesthesia of skin and scar neuromas. The nail first bend in the shape of "S" as a shape of clavicle to facilitate easy entry within narrow cavity and beyond fracture site. The nail will be used to create a path in the lateral end of the clavicle for subsequent easy access. The nail will be then passed from the medial side and across the reduced fracture into the lateral end of clavicle. The nail is cut underneath the skin to aid implant removal after complete union. Wound will be closed in layers and skin closure will be done. Sterile dressing will be done. [35]

Postoperatively the operated limb will be mobilized immediately on post-op day-1. IV antibiotics will be continued for 48 hrs and switched over to oral antibiotics on the 3rd day and continued till the 5th day.

Wound will be inspected on 3rd postoperative day and Sutures will be removed on 15th postoperative day.

Patient will be discharged on 2nd postoperative day but some patients will be discharged later due to

associated injuries and co morbidity. Rehabilitation will be started at end of 2 weeks. Gentle pendulum exercises to the shoulder will be allowed but abduction will be limited to 80 degrees. At 3 weeks active range of motion in all planes will be allowed. Regular follow up for every 4 week will be done. At each follow up patients will be assessed clinically, radiologically and the complications will be noted. Nail to be removed after 6 month. Local examinations of the affected clavicle for tenderness, instability, deformity and shoulder movements will be assessed. Functional assessment will be done by Constant Murley score

Fig. 1: Entry Point of TENS nailing



Figure 2: Pre-Operative X-Rays of a 55-year-old female patient with AO Type B1



Figure 3: Operative Instruments and Implant





Figure 4: (1 to 4): Intraoperative images showing entry of awl from medial end of clavicle ,with TENS nail attached to the T-handle passed through entry point and reduction of fracture achieved after manual shoulder movement. TENS nail passed beyond the fracture site and final anatomical reduction in a 55-year-old female with AO Type B1 Left Midshaft Clavicle fracture. Reduction forceps were used in some cases for fracture reduction.




Figure 5 : Post-Operative X-Rays



Figure 6: Pre operative and 5month follow up X-Rays of a -34 year-old male patient with AO Type B1 Left Midshaft Clavicle Fracture




Figure 7: Pre-Operative & 5 month follow up X-Rays of a 22-year-old male patient with AO Type B1 Left Midshaft clavicle fracture




Figure 8: Pre-Operative & 5 month follow up X-Rays of a 30-year-old female patient with AO Type B1 Left Midshaft clavicle fracture




Figure 9: Pre-Operative & 5 month follow up X-Rays of a 45-year-old male patient with AO Type B1 Right Midshaft clavicle fracture



Results:

In our study, we included 30 patients aged 18-50 years with closed midshaft clavicle fracture. 15 patients were males and 15 were females (Table 2).

Amongst them (12 right and 18 left side fracture) (Table 4), fracture result from road traffic accident, sport injury and fall. (Table 3).

There were 11 (type B1) and 19 (type B2) fractures. (Table 5). All patients were operated within 2 days following injury. The average operative time 30 min (range 25-60 min).

Among 30 patients we achieved closed reduction of fracture in 25 patients with lesser blood loss while in remaining 5 patients required open reduction of fracture site using bone holding and artery forcep. In all patient clinical union occurs at 5 weeks on an average (Range 4-6), and radiologically union occurs on an average 10 weeks. (Table 7) The patient were followed up post operatively and constant and murley score were calculated.

Constant Murley score at follow up

In the present study at follow up at 1 month 62.77 mean Constant and Murley score was observed, at 2 months 74.97 mean Constant and Murley score was observed, at 3 months 81.90 mean Constant and Murley score was observed and at 6 months 90.73 mean Constant and Murley score was observed. At follow up statistically significant difference was observed in Constant and Murley score. ($p < 0.0001$ ***)

Discussion

Clavicle fracture which are undisplaced, Midclavicular fractures can heal without the doctor conservatively, with the doctor, and despite the doctor!'.²⁴ The clavicle is the only long bone that is horizontal. The upper limb and the trunk are connected by this one and only bone strut. With concave forwards laterally and convex forwards medially, it has an S-shape. The bone is flat in the outer third, tubular in the middle, and serves as a transitional region between the two cross-sections. This renders the centre portion of the clavicle vulnerable, increasing the risk of fracture. The clavicle articulates medially with the sternum via the sternoclavicular joint and laterally with the scapula through the acromio-clavicular joint.

Adults frequently sustain clavicle fractures, which make up between 2.6% and 5% of all injuries.²⁵⁻²⁸ A fall on the outstretched hand or a direct hit to the anterior chest wall are the two possible mechanisms of injury.²⁹ Proximal, mid-shaft, and distal fractures are the three types of clavicle fractures. The mid-shaft is the most frequently fractured area of the clavicle, where the tiny cross section of the bone and the usual compressive stress delivered to the shoulder result in bony failure. Eighty percent of all clavicular fractures are mid-shaft fractures.³⁰

They show signs of displacement to some extent. However, 15–25% of all fractures are distal third fractures. Medial third fractures are the least common, making up only 0-5% of all clavicular fractures.^{31,32} A figure-of-eight bandage or an arm sling can be used as a conservative treatment for many of these mid-shaft fractures. According to recent data, more severe fracture types may actually have a higher likelihood of malunion or non-union.³³ Clavicle malunion has been demonstrated by several writers to be a unique clinical entity with distinctive patient symptomatology. Therefore, the purpose of the current study was to investigate the function of intramedullary nailing using Tens nail in cases of midshaft clavicle fracture.

Conclusion

- For midshaft clavicle fractures, TENS nailing guarantees high fracture union rates, superior radiological and functional results, and a prompt return to normal life activities.
- Minimally invasive surgery hardly 1-1.5 cm incision and requiring less operative time during surgery
- As indicated by higher Constant Murley ratings, it showed notable gains in functional recovery.

- Flexible intramedullary nailing can therefore be suggested as the preferred fixing technique for mid-shaft clavicle fractures because TENS also produced better cosmetic results, quicker healing durations, and early mobilization when compared to more conventional techniques like plating.

Authors Contribution

All authors participated in each and every aspect of this study, including design, data collection, data analysis, interpretation of data, patient contact and follow up, review and final version to be published in the journal.

Conflict of Interest

The authors confirm that they have no any conflict of interest related to research, authorship and publication of this article.

Funding

No funding from government, private or any other non profit institutions was used while doing this study.

Table 1: Age distribution

Age	No of cases	Percentage (%)
≤ 20	1	3.33
21–30	13	43.33
31–40	8	26.67
41–50	8	26.67
Total	30	100.00

Table 2: Gender distribution

Gender	No of cases	Percentage (%)
Male	15	50
Female	15	50
Total	30	100

Table 3: Mode of injury

Mode of injury	No of cases	Percentage (%)
----------------	-------------	----------------

Fall	10	33.33
RTA	15	50.00
Sports Injury	5	16.67
Total	30	100

Table 4: Side

Side	No of cases	Percentage (%)
Left	18	60.00
Right	12	40.00
Total	30	100.00

Table 5:AO Classification

AO Classification	No of cases	Percentage (%)
B1	11	36.67
B2	19	63.33
Total	30	100.00

Table 6:Associated injuries

AssociatedInjuries	No of cases	Percentage (%)
BBLeg#	2	6.67
I/LScapula spine#	1	3.33
No any	27	90.00
Total	30	100.00

Table 7:Time for union

Timeforunion(in weeks)	No of cases	Percentage (%)
≤ 6	5	16.67
7–9	18	60.00
10–12	7	23.33
Total	30	100.00

Table 8: Shoulder Movement

Shoulder movement (in degree)	No of cases	Percentage (%)
Flexion		
≤ 150	23	76.67
>150	7	23.33
Abduction		
≤ 150	18	60
>150	12	40
ER		
≤ 75	21	70
>75	9	30

Table 9:Constant and Murley score at follow up

Follow up	Mean±S.D
At1month	62.77±4.90
At2 months	74.97±2.63
At3 months	81.90±2.04
At6 months	90.73±2.23

Follow up	Mean±S.D	P-value
At1month	62.77±4.90	<0.0001
At6 months	90.73±2.23	

References:

1. Duan X, Zhong G, Cen S, Huang F, Xiang Z. Plating versus intramedullary pin or conservative treatment for midshaft fracture of clavicle: a meta-analysis of randomized controlled trials. *Journal of Shoulder and Elbow Surgery*. 2011 Sep 1;20(6):1008-15.
2. Frigg A, Rillmann P, Perren T, Gerber M, Ryf C. Intramedullary nailing of clavicular midshaft fractures with the titanium elastic nail: problems and complications. *The American Journal of Sports Medicine*. 2009 Feb;37(2):352-9.
3. Hill JM. Closed treatment of displaced middle-third fractures of the clavicle gives poor results. *The Journal of Bone & Joint Surgery British Volume*. 1997 Jul 1;79(4):537-8.
4. Khalil A. Intramedullary screw fixation for midshaft fractures of the clavicle. *International orthopaedics*. 2009 Oct;33:1421-4.
5. McKee MD, Pedersen EM, Jones C, Stephen DJ, Kreder HJ, Schemitsch EH, Wild LM, Potter J. Deficits following nonoperative treatment of displaced midshaft clavicular fractures. *JBJS*. 2006 Jan 1;88(1):35-40.
6. Mueller M, Rangger C, Striepens N, Burger C. Minimally invasive intramedullary nailing of midshaft clavicular fractures using titanium elastic nails. *Journal of Trauma and Acute Care Surgery*. 2008 Jun 1;64(6):1528-34.
7. Schuind F, Pay-Pay E, Andrianne Y, Donkerwolcke M, Rasquin C, Burny F. External fixation of the clavicle for fracture or non-union in adults. *JBJS*. 1988 Jun 1;70(5):692-5.
8. Zlowodzki M, Zelle BA, Cole PA, Jeray K, McKee MD. Treatment of acute midshaft clavicle fractures: systematic review of 2144 fractures: on behalf of the Evidence-Based Orthopaedic Trauma Working Group. *Journal of orthopaedic trauma*. 2005 Aug 1;19(7):504-7.
9. Grassi FA, Tajana MS, D'Angelo F. Management of midclavicular fractures: comparison between nonoperative treatment and open intramedullary fixation in 80 patients. *Journal of Trauma and Acute Care Surgery*. 2001 Jun 1;50(6):1096-100.
10. Leppilahti J, Jalovaara P. Migration of Kirschner wires following fixation of the clavicle-a report of 2 cases. *Acta Orthopaedica Scandinavica*. 1999 Jan 1;70(5):517-9.
11. Lyons FA, Rockwood Jr CA. Migration of pins used in operations on the shoulder. *JBJS*. 1990 Sep 1;72(8):1262-7.
12. Naidoo P. Migration of a Kirschner Wire from the clavicle into the abdominal aorta. *Archives of Emergency Medicine*. 1991 Dec;8(4):292.
13. Neer CS. Nonunion of the clavicle. *Journal of the American Medical Association*. 1960 Mar 5;172(10):1006-11.
14. Neviaser JS. The treatment of fractures of the clavicle. *Surgical Clinics of North America*. 1963 Dec 1;43(6):1555-63.
15. ROWE CR. 4 An Atlas of Anatomy and Treatment of Midclavicular Fractures. *Clinical Orthopaedics and Related Research (1976-2007)*. 1968 May 1;58:29-42.
16. Society CO. Nonoperative treatment compared with plate fixation of displaced midshaft clavicular fractures. A multicenter, randomized clinical trial.
17. Hill JM. Closed treatment of displaced middle-third fractures of the clavicle gives poor results. *The Journal of Bone & Joint Surgery British Volume*. 1997 Jul 1;79(4):537-8.
18. McKee MD, Pedersen EM, Jones C, Stephen DJ, Kreder HJ, Schemitsch EH, Wild LM, Potter J. Deficits following nonoperative treatment of displaced midshaft clavicular fractures. *JBJS*. 2006 Jan 1;88(1):35-40.
19. Society CO. Nonoperative treatment compared with plate fixation of displaced midshaft clavicular fractures. A multicenter, randomized clinical trial.
20. Bostman O, Manninen M, Pihlajamaki H. Complications of plate fixation in fresh displaced midclavicular fractures. *Journal of Trauma and Acute Care Surgery*. 1997 Nov 1;43(5):778-83.
21. Poigenfürst J, Rappold G, Fischer W. Plating of fresh clavicular fractures: results of 122 operations. *Injury*. 1992 Jan 1;23(4):237-41.
22. Kuner EH, Schlickewei W, Mydla F. Surgical therapy of clavicular fractures, indications, technic, results. *Hefte zur Unfallheilkunde*. 1982 Jan 1;160:76-83.
23. Bronz G, Heim D, Pusterla C, Heim U. Osteosynthesis of the clavicle (author's transl). *Unfallheilkunde*. 1981 Aug 1;84(8):319-25.
24. Freeland A. Unstable adult midclavicular fracture. *Orthopedics*. 1990 Nov 1;13(11):1279-81.

25. Bostman O, Manninen M, Pihlajamäki H. Complications of Wilk KE. The shoulder. Chapter-15, In Malone TR, McPoil TG, Nitz AJ, editors, Orthopaedics and sports physical therapy, 3rd edition, St.Louis : Mosby 1997; 401-409 plate fixation in fresh displaced midclavicular fractures. J Trauma 1997; 43: 778-83
26. Jubel A, Andermahr J, Schiffer G, Tsironis K, Rehm KE. Elastic stable intramedullary nailing of midclavicular fractures with a titanium nail. Clinical Orthopaedics and Related Research®. 2003 Mar 1;408:279-85.
27. Chu CM, Wang SJ, Lin LC. Fixation of mid-third clavicular fractures with knowles pins: 78 patients followed for 2-7 years. Acta Orthopaedica Scandinavica. 2002 Apr 1;73(2).
28. Jubel A, Andermahr J, Faymonville C, Binnebösel M, Prokop A, Rehm KE. Reconstruction of shoulder-girdle symmetry after midclavicular fractures. Stable, elastic intramedullary pinning versus rucksack bandage: Elastisch stabile intramedulläre Osteosynthese vs. Rucksackverband. Der Chirurg. 2002 Oct;73:978-81.
29. Hartmann F, Hessmann MH, Gercek E, Rommens PM. Elastic intramedullary nailing of midclavicular fractures. Acta chirurgica Belgica. 2008 Jan 1;108(4):428-32.
30. Wang YC, Fu YC, Chou SH, Liu PC, Tien YC, Lu CC. Titanium Elastic Nail versus plate fixation of displaced midshaft clavicle fractures: a retrospective comparison study. The Kaohsiung Journal of Medical Sciences. 2015 Sep 1;31(9):473-9.
31. Masatwar PV, Chishti SN, Singh SN, Maske RG, Soring D, Parija D. A prospective study of operative management of simple midshaft clavicular fracture with titanium elastic nail (TEN). International Journal of Orthopaedics. 2016;2(4):210-3.
32. Itagi PB, Pattanshetty AS. A comparative study of plate fixation versus titanium elastic nail in treatment of displaced midshaft clavicle fractures. International Journal of Orthopaedics. 2021;7(4):672-6.
33. Jones LD, Grammatopoulos G, Kambouroglou G. Titanium elastic nails, open reduction internal fixation and non-operative management for middle third clavicle fractures: a comparative study. Eur J Orthop Traumatol. 2014; 24(3):323-9.